



## TAKING DRUGS ON TIME



For HIV treatment to work well, you need to always take your pills at the right time, without missing any doses.

Anti-HIV drugs protect the body for several hours at a time.



## 05 IMPORTANT POINTS

- It's essential to take all your doses of HIV treatment at the right times and in the right amounts.
- Taking anti-HIV drugs regularly will mean that there is always enough of the drugs in your body. This will keep HIV under control.
- Not taking drugs as prescribed can result in HIV being harder to treat in the future.

## 06 NOTES

### What will happen if I do not take the drugs on time?

If you miss more than one dose a month, there is an increased risk that your treatment will stop working.

There will be more HIV in your body. This will weaken your immune system.

Some of the new HIV will be 'resistant' to anti-HIV drugs. In other words, the drugs won't work as well.

You may need to change the anti-HIV drugs you take. The next set of drugs may not be as easy to take and could have more side-effects.

gt grupo de trabajo sobre tratamientos del VIH  
ENTIDAD DECLARADA DE UTILIDAD PÚBLICA  
ONG DE DESARROLLO

### SUBVENCIONA



POR SOLIDARIDAD  
OTROS FINES DE INTERÉS SOCIAL



Secretaría del Plan Nacional sobre el Sida



Generalitat de Catalunya  
Agència de Salut Pública  
de Catalunya



Generalitat de Catalunya  
Departament  
de Justícia



Ajuntament de Barcelona  
ÀREA DE QUALITAT DE VIDA,  
IGUALTAT I ESPORTS



Diputació  
de Barcelona  
ÀREA DE BENESTAR SOCIAL

### COLABORA



ETIS  
EQUIP DE TREBALL SOBRE  
IMMIGRACIÓ I SALUT



¿TIENES DUDAS  
SOBRE EL TEMA?  
PREGÚNTANOS

Tel. 93 458 26 41  
consultas@gtt-vih.org